

The Stationary Measure of a 2-type Totally Asymmetric Exclusion Process

Omer Angel*

July 2004

Abstract

We give a combinatorial description of the stationary measure for a totally asymmetric exclusion process (TASEP) with second class particles, on either $\mathbb{Z}$ or on the cycle $\mathbb{Z}_N$. The measure is the image by a simple operation of the uniform measure on some larger finite state space. This reveals a combinatorial structure at work behind several results on the TASEP with second class particles.

1 Introduction

In the normal totally asymmetric simple exclusion process (TASEP), a number of particles occupy some distinct vertices of a graph, which in our case will be taken to be either $\mathbb{Z}$ or the cycle $\mathbb{Z}_N$. Each particle moves to the position to its right at rate one whenever that position is unoccupied. This gives a Markov chain with continuous time on the set of configurations — sets of locations for the particles. It is well known (and easy to prove) that if there are a particles on $\mathbb{Z}_N$, then the stationary distribution is uniform among all $\binom{N}{a}$ possible states. Since in the cyclic case the total number of particles is invariant, any other stationary measure is some linear combination of these uniform measures. Since the set of stationary measures is convex, from now on we focus only on the extremal stationary measures.

*Laboratoire de Math., Université Paris-Sud, Bat. 425, 91405 Orsay Cedex, France.
angel@math.ubc.ca

In the infinite case of $\mathbb{Z}$ there are two types of extremal stationary measures. There are the so called trivial stationary measures where the particles occupy all positions to the right of some point and no other positions. The non-trivial stationary measures are those where each vertex is occupied with some probability $p \in [0, 1]$ independently of all other vertices. Note that these measures are limits as $N \rightarrow \infty$ of the unique stationary measure on $\mathbb{Z}_N$ with $[pN]$ particles. For additional background on the TASEP and many references see [6, 7].

We are interested in the process with second class particles defined as follows (see e.g. [2]). The particles are classified as first class and second class particles. Sites are in one of three states: empty, or occupied by a single particle of one class or the other. As with the simple process, each particle jumps to the position to its right at rate 1 when that position is empty. Additionally, whenever a first class particle has a second class particle to its right, the two swap places at rate 1 (thus second class particles may move in both directions).

It is interesting at this point to note that the TASEP on a cycle is equivalent to the following shuffling method of cards. The cards are arranged in a cycle. Each consecutive pair is chosen at rate 1, and the pair is sorted with the larger card to the right. Assume there are a high cards, b medium cards and c low cards, and that cards within the same class are not distinguished from one another. The dynamics are exactly equivalent to those of the exclusion process with high cards corresponding to first class particles, medium cards to second class particles and low cards to empty positions (see [1] for an application). It is of course interesting to study the case where there are more than three types of cards (or particles). These cases seem significantly harder to analyze, though experimental studies of small cases, as well as the extremal case of N different particles do show interesting phenomena.

We give one final equivalent interpretation of the process, that may be related to the fact that adding further classes of particles breaks known techniques for working with the model. Non-empty sites of the graph are occupied by either a particle or an anti-particle. Each **edge** is chosen at rate 1. If a particle can move right across the edge to an empty spot it does. If an anti-particle can move left across the edge to an empty space it does. Finally, if there is a particle on the left and an anti-particle on the right, they both move, exchanging their positions, and otherwise nothing happens. Thus there are particles moving right and anti-particles moving left, but the movements

have rate 1 at each edge rather than each particle. This is equivalent to the previous form, with anti-particles representing empty spaces, and empty spaces representing the second class particles. When writing out states of the process, we will use 1's for particles, 0's for anti-particles and *'s for empty spaces. We will mostly refer to this interpretation of the process from now on, and use the term TASEP to refer to this process as well.

If the initial state is such that only two of the three types of positions appear (i.e. if there are no particles of one type or the other, or if all sites are occupied), then the process is simply the well understood exclusion process. However, the relation to the 1-type case is deeper than that. If an observer sees only the particles, and ignores the anti-particles, treating them as empty spaces, then he observes a regular exclusion process. Thus in the stationary distribution with a particles and b anti-particles, the marginal of the positions of particles is uniform over all $\binom{N}{a}$ sets. Similarly, the anti-particles form an exclusion process on their own (up to a reversal of the directions) and are therefore uniformly distributed over $\binom{N}{b}$ possible sets. Of course the two sets of locations are disjoint and hence dependent.

A similar statement holds in the infinite space case. In a stationary measure, if one type of particles are in a trivial state, then so must be the other, and we get the trivial stationary measures where a single segment of empty spaces has particles on its right and anti-particles on its left. Otherwise, the set of positions of particles have an i.i.d. product measure with some density, as do the anti-particles. Of course, the two marginals are not independent, nor is their joint distribution a product measure.

The main result of this paper is a combinatorial description of the stationary measures for the TASEP with three types of positions (empty and two types of particles). These results are then used to shed light on some known properties of the stationary measures. Duchi and Schaeffer [3] have found a similar relation, and use it to derive other results on the TASEP (primarily on a finite interval).

2 Definitions and Results

Definition 2.1. Two sets of positions S, T , of positions on $\mathbb{Z}$ or $\mathbb{Z}_N$, (not necessarily disjoint), are said to *collapse* to a state x of the TASEP if x is the result of the following (collapsing) procedure: Anti-particles are placed at the locations specified by T . Next, the locations in S are checked (in an

arbitrary order). If a location is empty, a particle is placed there. Otherwise a particle is placed in the nearest empty position **to the left** of the specified location.

In the case of the cycle $\mathbb{Z}_N$, we will only use the process for sets S, T with $|S| + |T| \leq n$, so that there is always an open position for every particle in S . In the case of $\mathbb{Z}$, if there are no open positions to the left of some element in S , we disregard that element.

The collapsing procedure defines a function from the product space (pairs of sets) to the space of states of the TASEP. The order at which the positions of S are used has no effect on the final resulting state x : The anti-particles in x are located exactly in the positions in T . A position a contains a particle in the resulting state x if and only if $a \notin T$ and there is some interval $I = [a, b]$ such that $|I \cap S| + |I \cap T| \geq |I|$. These statements may be used as an equivalent condition of collapsing, avoiding the possibly infinite algorithmic definition in the case of $\mathbb{Z}$.

Theorem 2.2. *The stationary measure for the exclusion process on $\mathbb{Z}_N$ with a particles and b anti-particles is the image by collapsing of the uniform measure on pairs of sub-sets S, T of the cycle of sizes a, b respectively.*

As a corollary, it follows that the least likely states for given population sizes are $1 \dots 1 * \dots * 0 \dots 0$ and its cyclic shifts, i.e. positions where sites of each type form a single interval, with positive particles to the left of the empty sites (and anti-particles to the right). These states have probability $1/\binom{N}{a}\binom{N}{b}$ each, while every other state has a probability that is some integer multiple of this probability. Section 6 contains some conjectures generalizing these facts to the case of more particle types.

As noted above, On $\mathbb{Z}$ the TASEP with second class particles has trivial stationary measures, which correspond to the states with particles to right of some point, and anti-particles to the left of some other point with empty spaces in between. In the notation of first and second class particles these are states with second class particles in some interval, first class particles to the right and empty spaces to the left.

Theorem 2.3. *The non-trivial extremal invariant measures for the exclusion process on $\mathbb{Z}$ are the image by collapsing of sets S, T where each $n \in S$ with probability p and $n \in T$ with probability q , all independently.*

Note that if $p + q \geq 1$, then a.s. all positions of $\mathbb{Z}$ will eventually have some particle on them, and then the resulting measure is one where each site has a particle with probability $1 - q$, and an anti-particle otherwise, independently of all other sites. As noted above, in the lack of empty spaces these are the (non-trivial extremal) invariant measures. If $p + q < 1$, then the resulting measure has particles with density p , anti-particles with density q and empty sites with density $1 - p - q$.

Theorems 2.2 and 2.3 can also be used to derive properties of the collapsing procedure. Since the distribution of the positions of particles in the stationary distribution is known:

Corollary 2.4. *Let S' be the set of locations of particles after collapsing independent S, T , which have either the uniform distribution over sets of a given size (in $\mathbb{Z}_N$) or i.i.d. product distributions (on $\mathbb{Z}$). Then S' also has a uniform or an i.i.d. product distribution.*

Furthermore, the TASEP itself has a natural symmetry of reversing the charge of the particles as well as the direction of the graph. It follows that a similar “dual” collapsing procedure where the particles are fixed and the anti-particles are moved forward to empty spaces also yields the same stationary measure.

In the next section we prove a combinatorial lemma that is closely related to stationarity of the collapsed uniform measures. Sections 4 and 5 contain the proof of Theorems 2.2, and 2.3 respectively. In Section 5 we also use the collapsing description of the stationary measures to shed new light on some of the results of [4]. Finally, Section 6 contains some open problems and conjectures regarding more general multi-type asymmetric exclusion processes.

3 Binary trees and dominated sequences

This section contains the combinatorial foundation for proving the stationarity of the collapsed uniform measure. The key result here is a bijection between binary trees and **pairs** of binary sequences, (i.e. sequences made up of 0's and 1's). The sequences are related to the TASEP since a binary sequence describes a segment in a state of the TASEP with no empty sites (which is one of the reasons we use the $\{0, *, 1\}$ notation).

Definition 3.1. Consider two finite binary sequences A, B of the same length n . We say that A *dominates* B and write $A \succcurlyeq B$ if it is possible to get from

A to B by moving 1's to the right. The *weight* of a binary sequence A is defined as the number of binary sequences dominated by it:

$$W(A) = |\{B : A \succcurlyeq B\}|.$$

In particular, it is necessary for $A \succcurlyeq B$ that both sequences have the same number of ones. Let the number of 1's in the first i digits of A (resp. B) be denoted by $a(i)$ (resp. $b(i)$). A condition equivalent to $A \succcurlyeq B$ is that $a(i) \geq b(i)$ for all i , and $a(n) = b(n)$. Thus for example, $W(1010) = 5$. If $x = 1 \dots 10 \dots 0$ has k ones followed by l zeroes, then $W(x) = \binom{k+l}{k}$. If the 1's and 0's were exchanged $W(x)$ would have been 1.

We use the following version of binary trees:

Definition 3.2. A *binary tree* is a rooted tree where each vertex including the root may have a left child, marked as left, and may have a right child, marked as right. A vertex may have either child, both, or neither.

Thus a tree is either the empty tree with only a root vertex, or it has a left sub-tree, a right sub-tree or both. Note that having an empty sub-tree on some side is different from not having a sub-tree on that side.

Next, we define recursively a function f mapping binary trees to binary sequences, as follows. The empty tree (with no edges) encodes the empty sequence. Otherwise,

$$f(T) = \begin{cases} f(L)0 & \text{if } T \text{ has only a left sub-tree } L, \\ 1f(R) & \text{if } T \text{ has only a right sub-tree } R, \\ f(L)01f(R) & \text{if } T \text{ has sub-trees } L, R, \end{cases}$$

where e.g. $f(L)01f(R)$ means a concatenation of $f(L)$, 01 and $f(R)$. In this way any binary sequence may be encoded by a binary tree, though generally the encoding is not unique. The length of the sequence is the number of edges of the tree, and the number of 1's is the total number of right children. The following combinatorial Lemma and immediate Corollary relate binary trees and dominated sequences:

Lemma 3.3. *There exists a bijection between binary trees T and pairs A, B of binary sequences such that $A \succcurlyeq B$ and $A = f(T)$.*

Corollary 3.4. *The number of binary trees that encode a given binary sequence A is $W(A)$.*

Proof of Lemma 3.3. First, we define in a similar fashion a second function g , that maps binary trees to binary sequences. The empty tree is mapped to the empty sequence, and otherwise,

$$g(T) = \begin{cases} 0g(L) & \text{if } T \text{ has only a left sub-tree } L, \\ 1g(R) & \text{if } T \text{ has only a right sub-tree } R, \\ 0g(L)1g(R) & \text{if } T \text{ has sub-trees } L, R, \end{cases}$$

We now show that mapping a tree T to a pair of sequences $A = f(T)$, $B = g(T)$ gives a bijection satisfying the Lemma's requirements.

The following facts are clear: for sequences A, B, C, D such that $A \succcurlyeq B$ and $C \succcurlyeq D$ we have $A0 \succcurlyeq 0B$ and $AC \succcurlyeq BD$. By induction, it follows that $f(T) \succcurlyeq g(T)$: If the root has only a right child, then $f(T) = 1f(R) \succcurlyeq 1g(R) = g(T)$. If the root has only a left child, then $f(T) = f(L)0 \succcurlyeq 0g(L) = g(T)$. If the root has two offspring, then concatenation of the previous two cases yields $f(T) \succcurlyeq g(T)$.

To see that this mapping is a bijection, we show how to (recursively) recover from sequences A, B a tree that is mapped to them, in such a way that at each step there is a unique possibility, so that the tree is unique.

Consider a binary tree T , with two sub-trees L, R and sequences $A = f(T) = f(L)01f(R)$ and $B = g(T) = 0g(L)1g(R)$. Recall that $a(i)$ and $b(i)$ count 1's in preambles of A and B . Since $f(L) \succcurlyeq g(L)$, we have that $a(i) \geq b(i+1)$ for $i \leq |L|$. However, $a(|L|+1) = b(|L|+1)$ is the number of right edges in L , and the next bit in B is a one. Thus the inequality $a(i) \geq b(i+1)$ fails for the first time for $i = |L|+1$. The following algorithm emerges: Given A, B find the first i for which $a(i) < b(i+1)$, and set $|L| = i - 1$. This generally identifies a unique representation of the sequences as $A = X01Y$ and $B = 0X'1Y'$ with $X \succcurlyeq X'$ and $Y \succcurlyeq Y'$. To reconstruct the unique tree T mapped to (A, B) , proceed recursively to identify L from X, X' and R from Y, Y' .

It remains to see that the cases where the above procedure fails to locate a representation as above correspond exactly to cases where T has only a left or only a right sub-tree. One possibility is that $a(0) = 0 < 1 = b(1)$, and then the above would give $|L| = -1$. In this case $A = 1Y$ and $B = 1Y'$ with $Y \succcurlyeq Y'$, so the unique tree mapped to A, B has only a right sub-tree R , where R is mapped to Y, Y' .

The other extreme is the case that $a(i) \geq b(i+1)$ for all $i < n$, suggesting $|R| = -1$. In this case we have $A = X0$ and $B = 0X'$ with $X \succcurlyeq X'$, and we

find a tree with only a left sub-tree. Thus in all cases the algorithm proceeds recursively to find a unique tree that is mapped to A, B . $\square$

Lemma 3.5. *For a binary sequence A ,*

$$W(A) = W(X)\mathbb{1}_{A=X0} + W(Y)\mathbb{1}_{A=1Y} + \sum_{X01Y=A} W(X)W(Y).$$

Thus if A ends with a 0 and equals $X0$, the RHS gets a contribution of $W(X)$ from the first term. Similarly, if A begins with a 1 there is a contribution from the second term. The sum in the RHS has a term in the sum for each representation of A as $X01Y$. For example, if $A = 1011010$ then we get

$$\begin{aligned} W(A) &= W(101101) + W(011010) + W(1)W(1010) + W(1011)W(0) \\ &= 7 + 9 + 1 \cdot 5 + 2 \cdot 1 = 23. \end{aligned}$$

Proof. By Corollary 3.4 it suffices to show that the RHS equals to the number of binary trees that encode A . This is done by induction. If $A = X0$ ends with a 0, then there are $W(X)$ trees encoding A with only a left sub-tree (which can be any tree encoding X). If $A = 1Y$ then similarly there are $W(Y)$ trees encoding A with only a right sub-tree. Finally, for any occurrence of 01 in A where $A = X01Y$ there are $W(X)$ possible left sub-trees and $W(Y)$ possible right sub-trees, giving a term in the sum. $\square$

4 Proof of Theorem 2.2

Consider a state x of the exclusion process on the cycle. How many pairs of sets S, T collapse to x ? Since the collapsing procedure begins by placing the anti-particles at T , the unique T is given by the set of positions of anti-particles in x . There may be a number of different sets S that (together with T) collapse to the state x . In order for the collapsing process to reach x it is necessary that S contains none of the empty positions of x (positions marked with $\star$'s). The empty positions in x break up the cycle into a number of segments each containing a sequence of particles (1's) and anti-particles (0's). Denote the binary segments of x by $A_1, \dots, A_l$.

During the collapsing procedure, if an element $p \in S$ results in a particle being placed in some position q to the left of p , there can be no empty position in the interval $[q, p]$, since otherwise the particle would have been placed there

instead. Thus the elements of S in each such binary segment must collapse into the positions marked for particles in that segment. It follows that for each binary segment A_i , the sequence having 1's at the elements of S in that segment is dominated by A_i , and so there are $W(A_i)$ possibilities for the intersection of S with that segment. The total number of possibilities for S is therefore $\prod W(A_i)$, and the collapsed uniform measure of the state x is

$$\mathbb{P}(x) = \frac{\prod W(A_i)}{\binom{N}{a} \binom{N}{b}}.$$

For example the cyclic state $*10**10100*0101$ may be reached from

$$W(10)W(\phi)W(10100)W(0101) = 2 \cdot 1 \cdot 9 \cdot 2 = 36$$

sets S and so its probability is $36/\binom{15}{5}\binom{15}{6}$.

To show that the collapsed uniform measure is stationary, place at each state x a mass $m(x) = \prod W(A_i)$ — a multiple of the collapsed uniform measure. Let the mass flow according to the transition kernel of the process (so if the process passes from x to y at rate r , mass flows from x to y at a rate of $r \cdot m(x)$). It suffices to show that the derivative of the mass at any state x is 0.

Denote $x \rightarrow_e y$ if an action (sorting particles) along an edge e leads from state x to state y . Since mass flowing from x to itself makes no difference, we only use this notation for $x \neq y$. Since the edge is determined uniquely by x and y , the edge subscript will usually be omitted. Each edge is used at rate 1, so the mass derivative is given by

$$\frac{d}{dt}m(x) = \sum_{y \rightarrow_e x} m(y) - \sum_{x \rightarrow_e z} m(x) = m(x) \sum_{y \rightarrow_e x} \left(\frac{m(y)}{m(x)} - \sum_{x \rightarrow_e z} 1 \right). \quad (1)$$

We now associate each term in each of the sums of (1) with one of the binary sequences A_i appearing in x in such a way that the terms associated with each sequence will cancel out, proving the Theorem. Since edges connecting two empty positions lead from x to itself and have been disregarded, each edge intersects exactly one of the binary segments of x , and the corresponding term is associated with that sequence.

The mass in the first sum — flowing into x — corresponds to edges with end-points marked '01', '*1', or '0*' in x . For each such edge we need to calculate the mass at the state resulting from 'unsorting' the edge. When

such an edge is unsorted, the resulting state y is very similar to x . Indeed, $\{B_i\}$ are the binary sequences appearing in y , then all but at one or two of them are equal to those in x .

- Consider first the case where $y \rightarrow_e x$ and the endpoints of e are marked '0*' in x and '*0' in y . In this case y has the same binary sequences as x except for two: $A_i = B_i 0$ and $B_{i+1} = 0 A_{i+1}$. Since $W(0A) = W(A)$, it follows that $\frac{m(y)}{m(x)} = \frac{W(B_i)}{W(A_i)}$, where B_i is A_i with a terminating 0 removed.
- Similarly, if $y \rightarrow_e x$ and the endpoints of e are marked '*1' in x and '1*' in y , In this case $A_i = 1 B_i$ and $B_{i-1} = A_{i-1} 1$. Since $W(A1) = W(A)$, we find that $\frac{m(y)}{m(x)} = \frac{W(B_i)}{W(A_i)}$, where B_i is A_i with an initial 1 removed.
- Finally, if e is marked with '01' in x , then splitting around that edge we have $A_i = X01Y$ and $B_i = X10Y$, with all other sequences beeing equal. Again, $\frac{m(y)}{m(x)} = \frac{W(B_i)}{W(A_i)}$.

Let us extend the $\rightarrow$ notation to binary sequences, so that $A \rightarrow B$ if it is possible to pass from A to B by either removing an initial 0, removing a terminating 1, or replacing a 10 by 01 somewhere in A . Consider the terms in the RHS of (1) that are associated with the binary sequence A_i of x . After substituting the above for $\frac{m(y)}{m(x)}$, these terms come to

$$m(x) \left(\sum_{B \rightarrow A_i} \frac{W(B)}{W(A_i)} - \sum_{A_i \rightarrow C} 1 \right),$$

and so it suffices to prove for an arbitrary sequence A that

$$\sum_{B \rightarrow A} W(B) = \sum_{A \rightarrow C} W(A). \quad (2)$$

Given a binary sequence A , associated terms in the RHS correspond to each occurrence of '10' in the sequence, as well as to an initial 0 if there is one and to terminating 1 if the sequence ends with a 1. The number of such terms is always one more than the number of times '01' appears in A .

The terms in the LHS, are $W(B)$ where B results from A either by replacing '01' by '10' at some place, or by removing an initial 1 or a terminating 0 if A has them. Consider a pair of sequences $B \rightarrow A$ where $A = X01Y$ and $B = X10Y$. Since $B \succ A$, B also dominates any sequence that A dominates.

The difference $W(B) - W(A)$ corresponds to sequences that B dominates and A does not. Such a sequence must be of the form $X'10Y'$, where $X \succcurlyeq X'$ and $Y \succcurlyeq Y'$, and so $W(B) = W(A) + W(X)W(Y)$.

Substituting this for $W(B)$ in (2) results in cancellation of all but one term on the RHS, and the resulting needed identity is exactly that given by Lemma 3.5.

5 The infinite setting

Proof of Theorem 2.3. It is known (see [5]) that there is a unique non-trivial stationary measure with marginals p, q for particles and anti-particles, and that those are all the non-trivial extremal stationary measures. It only needs to be shown that the collapsed i.i.d. measures are stationary and have the correct marginals.

The collapsed i.i.d. measure is the limit as $N \rightarrow \infty$ of the collapsed uniform measure on a cycle of length N with $[pN]$ particles and $[qN]$ anti-particles. Since in the finite case particles and anti-particles are uniformly distributed over all subsets of the appropriate size, in the limit they have densities p and q respectively. Since correlations in this measure decay exponentially in the distance, it follows that the limit is stationary. $\square$

The collapsing procedure sheds new light some known results. First, the fact that second class particles induce a factoring of the stationary distribution (see [2]): given that 0 is an empty position, the state in $\mathbb{Z}^+$ and in $\mathbb{Z}^-$ are independent. With the collapsing procedure, the state in $\mathbb{Z}^+$ depends only on the positive elements of S, T . Conditioned on the event that 0 remains empty, no particle crosses from $\mathbb{Z}^+$ to $\mathbb{Z}^-$, and so the state on $\mathbb{Z}^-$ is determined by the negative elements of S, T and is therefore independent of the state on $\mathbb{Z}^+$.

Next, consider the relation demonstrated in [8, 4] to a certain biased random walk on $\mathbb{Z}$. For marginals p for particles and q for anti-particles, the corresponding random walk has steps of $-1, 0, 1$ with step distribution given by $\mathbb{P}(X = 1) = (1 - p)(1 - q)$, and $\mathbb{P}(X = -1) = pq$. When $p + q < 1$ we have $\mathbb{E}X = 1 - p - q > 0$.

The random walk is very naturally coupled with the pair S, T of i.i.d. subsets of $\mathbb{Z}$ with densities p, q , and is given by $Z_n = |T^c \cap [-n, 0)| - |S \cap [-n, 0)|$ (note that the sets are explored backwards), which clearly has the

above step distribution. Lemma 2.5 of [4] says that the distribution of the distance between second class particles is the same as the hitting time of 1 by the random walk. In our notation this is a statement regarding the distance between empty positions. Given S, T such that there is no particle at 0, the next hole to the left of 0 is at $-n$ exactly when the corresponding Z hits 1 at time n .

The following is standard and easily seen: The stationary measures seen from a single second class particles are derived from the stationary measures with anti-particles, by the following rule. Condition on having an empty position at 0, and place a second class particle there. All particles are first class particles, and any empty spot to the right of 0 is also filled with a first class particle. All other positions become empty. By selecting the densities of particles and anti-particles the asymptotic densities to either side can be controlled.

Extend the definition of the random walk to negative indices by $Z_{-n} = -|T^c \cap [0, n)| + |S \cap [0, n)|$. Theorem 5 of [4] states that the stationary measure for the process seen from a single second class particle with given asymptotic densities is very close to a product of independent i.i.d. measures on $\mathbb{Z}^+$ and $\mathbb{Z}^-$ with the given densities. Formally, the stationary measure and the product measure can be coupled so that they are exponentially unlikely to differ in many positions.

This again follows from the collapsing procedure, since the final location of a particle is exponentially unlikely to be far from the position given by S .

6 Open problems

The following conjectures are a generalization of our results to more classes of particles, both in qualitative and quantitative terms.

Conjecture 1. *In a cycle with particles of a number of classes, in the stationary distribution, the least likely states are those where if the cycle is cut at some point the particles are arranged in reversed order of speed.*

Let x be one of the above states (there are N of them).

Conjecture 2. *In the stationary distribution, $\mathbb{P}(x) = \prod \binom{N}{s_i}^{-1}$, where n is the cycle length and s_i is the number of particles of class at least i .*

Conjecture 3. *In the stationary distribution, the probability of any other state is an integral multiple of $\mathbb{P}(x)$.*

Finally, is there a useful generalization of the collapsing procedure for processes with more classes of particles? The obvious generalizations of repeated collapsing do not appear to give the correct stationary distribution. The smallest case where they fail is that of a cycle of length 4 with all different particles (or 3 particles and an empty space). It turns out that $\mu(1324) \neq \mu(1423)$ under the stationary measure μ .

References

- [1] I. Benjamini, N. Berger, C. Hoffman, and E. Mossel. Mixing times of the biased card shuffling and the asymmetric exclusion process. To appear in Trans. of AMS., 2004.
- [2] B. Derrida, S. Janowsky, J. Lebowitz, and E. Speer. Exact solution of the totally asymmetric simple exclusion process: shock profiles. *J. Stat. Phys.*, 73:833–874, 1993.
- [3] E. Duchi and G. Schaeffer. A combinatorial approach to jumping particles. To appear, J. Comb. The. A., 2004.
- [4] P. Ferrari, L. Fontes, and Y. Kohayakawa. Invariant measures for a two species asymmetric process. *J. Stat. Phys.*, 76(5):1153–1177, 1994.
- [5] P. Ferrari, C. Kipnis, and E. Saada. Microscopic structure of traveling waves in the asymmetric simple exclusion process. *Ann. Prob.*, 19:226–244, 1991.
- [6] T. Liggett. *Interacting Particle Systems*. Fundamental Principles of Mathematical Sciences, 276. Springer-Verlag, New York, 1985.
- [7] T. Liggett. *Stochastic Interacting Systems: Contact, Voter and Exclusion Processes*. Fundamental Principles of Mathematical Sciences, 324. Springer-Verlag, New York, 1999.
- [8] Speer. The two species asymmetric simple exclusion process. In A. V. M. Fannes, C. Maes, editor, *Micro, Meso, and Macroscopic Approaches in Physics, Proceedings of the NATO AR Workshop "On Three Levels"*, Leuven, July 1993 1994.